

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

|  |    |   |
|--|----|---|
| (51) International Patent Classification <sup>4</sup> :<br><br>C12Q 1/68   | A1 | (11) International Publication Number: <b>WO 89/10977</b><br><br>(43) International Publication Date: 16 November 1989 (16.11.89)   |
| <p>(21) International Application Number: PCT/GB89/00460</p> <p>(22) International Filing Date: 2 May 1989 (02.05.89)</p> <p>(30) Priority data:<br/>8810400.5                      3 May 1988 (03.05.88)                      GB</p> <p>(71) Applicant (for all designated States except US): ISIS INNOVATION LIMITED [GB/GB]; 2 South Parks Road, Oxford OX1 3UB (GB).</p> <p>(72) Inventor; and<br/>(75) Inventor/Applicant (for US only) : SOUTHERN, Edwin [GB/GB]; 30 Staverton Road, Oxford OX2 6XJ (GB).</p> <p>(74) Agent: PENNANT, Pyers; Stevens, Hewlett &amp; Perkins, 5 Quality Court, Chancery Lane, London WC2A 1HZ (GB).</p>   |    | <p>(81) Designated States: AT (European patent), BE (European patent), CH (European patent), DE (European patent), FR (European patent), GB (European patent), IT (European patent), JP, LU (European patent), NL (European patent), SE (European patent), US.</p> <p><b>Published</b><br/><i>With international search report.</i></p> |
| <p>(54) Title: ANALYSING POLYNUCLEOTIDE SEQUENCES</p> <p>(57) Abstract</p> <p>This invention provides apparatus and method for analysing a polynucleotide sequence, either an unknown sequence or a known sequence. A support, e.g. a glass plate, carries an array of the whole or a chosen part of a complete set of oligonucleotides which are capable of taking part in hybridization reactions. The array may comprise one or more pairs of oligonucleotides. The polynucleotide sequence, or fragments thereof, are labelled and applied to the array under hybridizing conditions. Applications include analysis of known point mutations, genomic fingerprinting, linkage analysis, characterization of mRNAs, mRNA populations, and sequence determination.</p> |    |   |

***FOR THE PURPOSES OF INFORMATION ONLY***

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

|    |                              |    |  |    |                          |
|----|------------------------------|----|--|----|--------------------------|
| AT | Austria                      | FI | Finland                                  | ML | Mali                     |
| AU | Australia                    | FR | France                                   | MR | Mauritania               |
| BB | Barbados                     | GA | Gabon                                    | MW | Malawi                   |
| BE | Belgium                      | GB | United Kingdom                           | NL | Netherlands              |
| BF | Burkina Faso                 | HU | Hungary                                  | NO | Norway                   |
| BG | Bulgaria                     | IT | Italy                                    | RO | Romania                  |
| BJ | Benin                        | JP | Japan                                    | SD | Sudan                    |
| BR | Brazil                       | KP | Democratic People's Republic<br>of Korea | SE | Sweden                   |
| CF | Central African Republic     |    |  | SN | Senegal                  |
| CG | Congo                        | KR | Republic of Korea                        | SU | Soviet Union             |
| CH | Switzerland                  | LI | Liechtenstein                            | TD | Chad                     |
| CM | Cameroon                     | LK | Sri Lanka                                | TG | Togo                     |
| DE | Germany, Federal Republic of | LU | Luxembourg                               | US | United States of America |
| DK | Denmark                      | MC | Monaco                                   |    |                          |
| ES | Spain                        | MG | Madagascar                               |    |                          |

ANALYSING POLYNUCLEOTIDE SEQUENCES

## 1. INTRODUCTION

Three methods dominate molecular analysis of nucleic acid sequences: gel electrophoresis of restriction fragments, molecular hybridisation, and the rapid DNA sequencing methods. These three methods have a very wide range of applications in biology, both in basic studies, and in the applied areas of the subject such as medicine and agriculture. Some idea of the scale on which the methods are now used is given by the rate of accumulation of DNA sequences, which is now well over one million base pairs a year. However, powerful as they are, they have their limitations. The restriction fragment and hybridisation methods give a coarse analysis of an extensive region, but are rapid; sequence analysis gives the ultimate resolution, but it is slow, analysing only a short stretch at a time. There is a need for methods which are faster than the present methods, and in particular for methods which cover a large amount of sequence in each analysis.

This invention provides a new approach which produces both a fingerprint and a partial or complete sequence in a single analysis, and may be used directly with complex DNAs and populations of RNA without the need for cloning.

In one aspect the invention provides apparatus for analysing a polynucleotide sequence, comprising a support and attached to a surface thereof an array of the whole or a chosen part of a complete set of oligonucleotides of chosen lengths, the oligonucleotides being capable of taking part in hybridisation reactions. For studying differences between polynucleotide sequences, the invention provides in another aspect apparatus comprising a support and attached to a surface thereof an array of the whole or a chosen part of a complete set of oligonucleotides of chosen lengths comprising the polynucleotide sequences, the oligo-

nucleotides being capable of taking part in hybridisation reactions.

5 In another aspect, the invention provides a method of analysing a polynucleotide sequence, by the use of a support to the surface of which is attached an array of the whole or a chosen part of a complete set of oligo-  
nucleotides of chosen lengths, which method comprises  
labelling the polynucleotide sequence or fragments  
thereof to form labelled material, applying the  
10 labelled material under hybridisation conditions to the array, and observing the location of the label on the surface associated with particular members of the set of oligonucleotides.

15 The idea of the invention is thus to provide a structured array of the whole or a chosen part of a complete set of oligonucleotides of one or several chosen lengths. The array, which may be laid out on a supporting film or glass plate, forms the target for a hybridisation reaction. The chosen conditions of  
20 hybridisation and the length of the oligonucleotides must at all events be sufficient for the available equipment to be able to discriminate between exactly matched and mismatched oligonucleotides. In the hybridisation reaction, the array is explored by a  
25 labelled probe, which may comprise oligomers of the chosen length or longer polynucleotide sequences or fragments, and whose nature depends on the particular application. For example, the probe may comprise  
30 labelled sequences amplified from genomic DNA by the polymerase chain reaction, or a mRNA population, or a complete set of oligonucleotides from a complex sequence such as an entire genome. The end result is a set of filled cells corresponding to the  
oligonucleotides present in the analysed sequence, and  
35 a set of "empty" sites corresponding to the sequences

which are absent in the analysed sequence. The pattern produces a fingerprint representing all of the sequence analysed. In addition, it is possible to assemble most or all of the sequence analysed if an oligonucleotide length is chosen such that most or all oligonucleotide sequences occur only once.

The number, the length and the sequences of the oligonucleotides present in the array "lookup table" also depend on the application. The array may include all possible oligonucleotides of the chosen length, as would be required if there was no sequence information on the sequence to be analysed. In this case, the preferred length of oligonucleotide used depends on the length of the sequence to be analysed, and is such that there is likely to be only one copy of any particular oligomer in the sequence to be analysed. Such arrays are large. If there is any information available on the sequence to be analysed, the array may be a selected subset. For the analysis of a sequence which is known, the size of the array is of the same order as length of the sequence, and for many applications, such as the analysis of a gene for mutations, it can be quite small. These factors are discussed in detail in what follows.

## 2. OLIGONUCLEOTIDES AS SEQUENCE PROBES

Oligonucleotides form base paired duplexes with oligonucleotides which have the complementary base sequence. The stability of the duplex is dependent on the length of the oligonucleotides and on base composition. Effects of base composition on duplex stability can be greatly reduced by the presence of high concentrations of quaternary or tertiary amines. However, there is a strong effect of mismatches in the oligonucleotides duplex on the thermal stability of the hybrid, and it is this which makes the technique of

hybridisation with oligonucleotides such a powerful method for the analysis of mutations, and for the selection of specific sequences for amplification by DNA polymerase chain reaction. The position of the mismatch affects the degree of destabilisation. Mismatches in the centre of the duplex may cause a lowering of the  $T_m$  by  $10^{\circ}\text{C}$  compared with  $1^{\circ}\text{C}$  for a terminal mismatch. There is then a range of discriminating power depending on the position of mismatch, which has implications for the method described here. There are ways of improving the discriminating power, for example by carrying out hybridisation close to the  $T_m$  of the duplex to reduce the rate of formation of mismatched duplexes, and by increasing the length of oligonucleotide beyond what is required for unique representation. A way of doing this systematically is discussed.

### 3. ANALYSIS OF A PREDETERMINED SEQUENCE

One of the most powerful uses of oligonucleotide probes has been in the detection of single base changes in human genes. The first example was the detection of the single base change in the betaglobin gene which leads to sickle cell disease. There is a need to extend this approach to genes in which there may be a number of different mutations leading to the same phenotype, for example the DMD gene and the HPRT gene, and to find an efficient way of scanning the human genome for mutations in regions which have been shown by linkage analysis to contain a disease locus for example Huntington's disease and Cystic Fibrosis. Any known sequence can be represented completely as a set of overlapping oligonucleotides. The size of the set is  $N - s + 1 \approx N$ , where  $N$  is the length of the sequence and  $s$  is the length of an oligomer. A gene of 1 kb for example, may be divided into an overlapping set of

- 5 -

around one thousand oligonucleotides of any chosen length. An array constructed with each of these oligonucleotides in a separate cell can be used as a multiple hybridisation probe to examine the homologous sequence in any context, a single-copy gene in the human genome or a messenger RNA among a mixed RNA population, for example. The lengths may be chosen such that there is only a small probability that any oligomer in the sequence is represented elsewhere in the sequence to be analysed. This can be estimated from the expression given in the section discussing statistics below. For a less complete analysis it would be possible to reduce the size of the array e.g. by a factor of up to 5 by representing the sequence in a partly or non-overlapping set. The advantage of using a completely overlapping set is that it provides a more precise location of any sequence difference, as the mismatch will scan in s consecutive oligonucleotides.

#### 4. ANALYSIS OF AN UNDETERMINED SEQUENCE

The genomes of all free living organisms are larger than a million base pairs and none has yet been sequenced completely. Restriction site mapping reveals only a small part of the sequence, and can detect only a small portion of mutations when used to compare two genomes. More efficient methods for analysing complex sequences are needed to bring the full power of molecular genetics to bear on the many biological problems for which there is no direct access to the gene or genes involved. In many cases, the full sequence of the nucleic acids need not be determined; the important sequences are those which differ between two nucleic acids. To give three examples: the DNA sequences which are different between a wild type organism and one which carries a mutant can lead the way to isolation of the relevant gene; similarly, the sequence differences between a cancer cell and its

- 6 -

normal counterpart can reveal the cause of transformation; and the RNA sequences which differ between two cell types point to the functions which distinguish them. These problems can be opened to molecular analysis by a method which identifies sequence differences. Using the approach outlined here, such differences can be revealed by hybridising the two nucleic acids, for example the genomic DNA of the two genotypes, or the mRNA populations of two cell types to an array of oligonucleotides which represent all possible sequences. Positions in the array which are occupied by one sequence but not by the other show differences in two sequences. This gives the sequence information needed to synthesise probes which can then be used to isolate clones of the sequence involved.

#### 4.1 ASSEMBLING THE SEQUENCE INFORMATION

Sequences can be reconstructed by examining the result of hybridisation to an array. Any oligonucleotide of length  $s$  from within a long sequence, overlaps with two others over a length  $s-1$ . Starting from each positive oligonucleotide, the array may be examined for the four oligonucleotides to the left and the four to the right that can overlap with a one base displacement. If only one of these four oligonucleotides is found to be positive to the right, then the overlap and the additional base to the right determine  $s$  bases in the unknown sequence. The process is repeated in both directions, seeking unique matches with other positive oligonucleotides in the array. Each unique match adds a base to the reconstructed sequence.

#### 4.2 SOME STATISTICS

Any sequence of length  $N$  can be broken down to a set of  $\sim N$  overlapping sequences  $s$  base pairs in length. (For double stranded nucleic acids, the sequence complexity of a sequence of  $N$  base pairs is  $2N$ , because the two strands have different sequences,



- 7 -

but for the present purpose, this factor of two is not significant). For oligonucleotides of length  $s$ , there are  $4^s$  different sequence combinations. How big should  $s$  be to ensure that most oligonucleotides will be represented only once in the sequence to be analysed, of complexity  $N$ ? For a random sequence the expected number of  $s$ -mers which will be present in more than one copy is

$$\mu_{>1} \approx 4^s(1 - e^{-\lambda(1+\lambda)})$$

10

where

$$\lambda = (N - s + 1)/4^s$$

For practical reasons it is also useful to know how many sequences are related to any given  $s$ -mer by a single base change. Each position can be substituted by one of three bases, there are therefore  $3s$  sequences related to an individual  $s$ -mer by a single base change, and the probability that any  $s$ -mer in a sequence of  $N$  bases is related to any other  $s$ -mer in that sequence allowing one substitution is  $3s \times N/4^s$ . The relative signals of matched and mismatched sequences will then depend on how good the hybridisation conditions are in distinguishing a perfect match from one which differ by a single base. (If  $4^s$  is an order of magnitude greater than  $N$ , there should only be a few,  $3s/10$ , related to any oligonucleotide by one base change.) The indications are that the yield of hybrid from the mismatched sequence is a fraction of that formed by the perfect duplex.

30

For what follows, it is assumed that conditions can be found which allow oligonucleotides which have complements in the probe to be distinguished from those which do not.

#### 4.3 ARRAY FORMAT, CONSTRUCTION AND SIZE

35

To form an idea of the scale of the arrays needed

- 8 -

to analyse sequences of different complexity it is convenient to think of the array as a square matrix. All sequences of a given length can be represented just once in a matrix constructed by drawing four rows  
 5 representing the four bases, followed by four similar columns. This produces a 4 x 4 matrix in which each of the 16 squares represents one of the 16 doublets. Four similar matrices, but one quarter the size, are then drawn within each of the original squares. This  
 10 produces a 16 x 16 matrix containing all 256 tetra-nucleotide sequences. Repeating this process produces a matrix of any chosen depth, s, with a number of cells equal to  $4^s$ . As discussed above, the choice of s is of great importance, as it determines the complexity of  
 15 the sequence representation. As discussed below, s also determines the size of the matrix constructed, which must be very big for complex genomes. Finally, the length of the oligonucleotides determines the hybridisation conditions and their discriminating power  
 20 as hybridisation probes.

|    |    |                      | Side of Matrix    | Number of                          |
|----|----|----------------------|-------------------|------------------------------------|
|    | s  | $4^s$                | Genomes           | (pixel=100 $\mu$ m) Sheets of film |
|    | 8  | 65536                | $4^s \times 10^6$ |                                    |
|    | 9  | 262144               |                   |                                    |
| 25 | 10 | $1.0 \times 10^6$    | cosmid            | 100 $\mu$ m 1                      |
|    | 11 | $4.2 \times 10^6$    |                   |                                    |
|    | 12 | $1.7 \times 10^7$    |                   |                                    |
|    | 13 | $6.7 \times 10^7$    | E.coli            |                                    |
|    | 14 | $2.6 \times 10^8$    | yeast             | 1.6 $\mu$ m 9                      |
| 30 | 15 | $1.1 \times 10^9$    |                   |                                    |
|    | 16 | $4.2 \times 10^9$    |                   |                                    |
|    | 17 | $1.7 \times 10^{10}$ |                   |                                    |
|    | 18 | $6.7 \times 10^{10}$ | human             | 25 $\mu$ m 2,500                   |
|    | 19 | $2.7 \times 10^{11}$ |                   |                                    |
| 35 | 20 | $1.1 \times 10^{12}$ |                   | 100 $\mu$ m                        |

- 9 -

The table shows the expected scale of the arrays needed to perform the first analysis of a few genomes. The examples were chosen because they are genomes which have either been sequenced by conventional procedures - the cosmid scale -, are in the process of being sequenced - the E. coli scale -, or for which there has been considerable discussion of the magnitude of the problem - the human scale. the table shows that the expected scale of the matrix approach is only a small fraction of the conventional approach. This is readily seen in the area of X-ray film that would be consumed. It is also evident that the time taken for the analysis would be only a small fraction of that needed for gel methods. The "Genomes" column shows the length of random sequence which would fill about 5% of cells in the matrix. This has been determined to be the optimum condition for the first step in the sequencing strategy discussed below. At this size, a high proportion of the positive signals would represent single occurrences of each oligomer, the conditions needed to compare two genomes for sequence differences.

#### 5. REFINEMENT OF AN INCOMPLETE SEQUENCE

Reconstruction of a complex sequence produces a result in which the reconstructed sequence is interrupted at any point where an oligomer that is repeated in the sequence occurs. Some repeats are present as components of long repeating structures which form part of the structural organisation of the DNA, dispersed and tandem repeats in human DNA for example. But when the length of oligonucleotide used in the matrix is smaller than that needed to give totally unique sequence representation, repeats occur by chance. Such repeats are likely to be isolated. That is, the sequences surrounding the repeated oligomers are unrelated to each other. The gaps caused

- 10 -

by these repeats can be removed by extending the sequence to longer oligomers. In principle, those sequences shown to be repeated by the first analysis, using an array representation of all possible oligomers, could be resynthesised with an extension at each end. For each repeated oligomer, there would be  $4 \times 4 = 16$  oligomers in the new matrix. The hybridisation analysis would now be repeated until the sequence was complete. In practice, because the results of a positive signal in the hybridisation may be ambiguous, it may be better to adopt a refinement of the first result by extending all sequences which did not give a clear negative result in the first analysis. An advantage of this approach is that extending the sequence brings mismatches which are close to the ends in the shorter oligomer, closer to the centre in the extended oligomer, increasing the discriminatory power of duplex formation.

#### 5.1 A HYPOTHETICAL ANALYSIS OF THE SEQUENCE OF BACTERIOPHAGE $\lambda$ DNA

Lambda phage DNA is 48,502 base pairs long. Its sequence has been completely determined, we have treated one strand of this as a test case in a computer simulation of the analysis. The table shows that the appropriate size of oligomer to use for a sequence of this complexity is the 10-mer. With a matrix of 10-mers, the size was 1024 lines square. After "hybridisation" of the lambda 10-mers in the computer, 46,377 cells were positive, 1957 had double occurrences, 75 triple occurrences, and three quadruple occurrences. These 46,377 positive cells represented known sequences, determined from their position in the matrix. Each was extended by four x one base at the 3' end and four x one base at the 5', end to give  $16 \times 46,377 = 742,032$  cells. This extended set reduced the

- 11 -

number of double occurrences to 161, a further 16-fold extension brought the number down to 10, and one more provided a completely overlapped result. Of course, the same end result of a fully overlapped sequence could be achieved starting with a  $4^{16}$  matrix, but the matrix would be 4096 times bigger than the matrix needed to represent all 10-mers, and most of the sequence represented on it would be redundant.

## 5.2 LAYING DOWN THE MATRIX

The method described here envisages that the matrix will be produced by synthesising oligonucleotides in the cells of an array by laying down the precursors for the four bases in a predetermined pattern, an example of which is described above. Automatic equipment for applying the precursors has yet to be developed, but there are obvious possibilities; it should not be difficult to adapt a pen plotter or other computer-controlled printing device to the purpose. The smaller the pixel size of the array the better, as complex genomes need very large numbers of cells. However, there are limits to how small these can be made. 100 microns would be a fairly comfortable upper limit, but could probably not be achieved on paper for reasons of texture and diffusion. On a smooth impermeable surface, such as glass, it may be possible to achieve a resolution of around 10 microns, for example by using a laser typesetter to preform a solvent repellant grid, and building the oligonucleotides in the exposed regions. One attractive possibility, which allows adaptation of present techniques of oligonucleotide synthesis, is to sinter microporous glass in microscopic patches onto the surface of a glass plate. Laying down very large number of lines or dots could take a long time, if the printing mechanism were slow. However, a low cost ink-

- 12 -

jet printer can print at speeds of about 10,000 spots per second. With this sort of speed,  $10^8$  spots could be printed in about three hours.

5

### 5.3 OLIGONUCLEOTIDE SYNTHESIS

There are several methods of synthesising oligonucleotides. Most methods in current use attach the nucleotides to a solid support of controlled pore size glass (CPG) and are suitable for adaptation to synthesis on a glass surface. Although we know of no description of the direct use of oligonucleotides as hybridisation probes while still attached to the matrix on which they were synthesised, there are reports of the use of oligonucleotides as hybridisation probes on solid supports to which they were attached after synthesis. PCT Application WO 85/01051 describes a method for synthesising oligonucleotides tethered to a CPG column. In an experiment performed by us, CPG was used as the support in an Applied Bio-systems oligonucleotide synthesiser to synthesise a 13-mer complementary to the left hand cos site of phage lambda. The coupling steps were all close to theoretical yield. The first base was stably attached to the support medium through all the synthesis and deprotection steps by a covalent link.

10

15

20

25

### 6. PROBES, HYBRIDISATION AND DETECTION

The yield of oligonucleotides synthesised on microporous glass is about 30  $\mu\text{mol/g}$ . A patch of this material 1 micron thick by 10 microns square would hold  $\sim 3 \times 10^{-12}$   $\mu\text{mol}$ , equivalent to about 2 g of human

30

35

- 13 -

DNA. The hybridisation reaction could therefore be carried out with a very large excess of the bound oligonucleotides over that in the probe. So it should be possible to design a system capable of distinguishing between hybridisation involving single and multiple occurrences of the probe sequence, as yield will be proportional to concentration at all stages in the reaction.

The polynucleotide sequence to be analysed may be of DNA or RNA. To prepare the probe, the polynucleotide may be degraded to form fragments. Preferably it is degraded by a method which is as random as possible, to an average length around the chosen length  $s$  of the oligonucleotides on the support, and oligomers of exact length  $s$  selected by electrophoresis on a sequencing gel. The probe is then labelled. For example, oligonucleotides of length  $s$  may be end labelled. If labelled with  $^{32}\text{P}$ , the radioactive yield of any individual  $s$ -mer even from total human DNA could be more than  $10^4$  dpm/mg of total DNA. For detection, only a small fraction of this is needed in a patch 10-100 microns square. This allows hybridisation conditions to be chosen to be close to the  $T_m$  of duplexes, which decreases the yield of hybrid and decreases the rate of formation, but increases the discriminating power. Since the bound oligonucleotide is in excess, signal need not be a problem even working close to equilibrium.

Hybridisation conditions can be chosen to be those known to be suitable in standard procedures used to hybridise to filters, but establishing optimum conditions is important. In particular, temperature needs to be controlled closely, preferably to better than  $\pm 0.5^\circ\text{C}$ . Particularly when the chosen length of the oligonucleotide is small, the analysis needs to be

- 14 -

able to distinguish between slight differences of rate and/or extent of hybridisation. The equipment may need to be programmed for differences in base composition between different oligonucleotides. In constructing  
5 the array, it may be preferable to partition this into sub-matrices with similar base compositions. This may make it easier to define the  $T_m$  which may differ slightly according to the base composition.

Autoradiography, especially with  $^{32}P$  causes image  
10 degradation which may be a limiting factor determining resolution; the limit for silver halide films is around 25 microns. Obviously some direct detection system would be better. Fluorescent probes are envisaged; given the high concentration of the target  
15 oligonucleotides, the low sensitivity of fluorescence may not be a problem.

We have considerable experience of scanning autoradiographic images with a digitising scanner. Our present design is capable of resolution down to 25  
20 microns, which could readily be extended down to less than present application, depending on the quality of the hybridisation reaction, and how good it is at distinguishing absence of a sequence from the presence of one or more. Devices for measuring astronomical  
25 plates have an accuracy around 1  $\mu$ . Scan speeds are such that a matrix of several million cells can be scanned in a few minutes. Software for the analysis of the data is straight-forward, though the large data sets need a fast computer.

30 Experiments presented below demonstrate the feasibility of the claims.

Commercially available microscope slides (BDH Super Premium 76 x 26 x 1 mm) were used as supports. These were derivatised with a long aliphatic linker  
35 that can withstand the conditions used for the



- 15 -

deprotection of the aromatic heterocyclic bases, i.e. 30%  $\text{NH}_3$  at  $55^\circ$  for 10 hours. The linker, bearing a hydroxyl group which serves as a starting point for the subsequent oligonucleotide, is synthesised in two steps. The slides are first treated with a 25% solution of 3-glycidoxypropyltriethoxysilane in xylene containing several drops of Hunig's base as a catalyst. The reaction is carried out in a staining jar, fitted with a drying tube, for 20 hours at  $90^\circ\text{C}$ . The slides are washed with  $\text{MeOH}$ ,  $\text{Et}_2\text{O}$  and air dried. Then neat hexaethylene glycol and a trace amount of conc. sulphuric acid are added and the mixture kept at  $80^\circ$  for 20 hours. The slides are washed with  $\text{MeOH}$ ,  $\text{Et}_2\text{O}$ , air dried and stored desiccated at  $-20^\circ$  until use. This preparative technique is described in British Patent Application 8822228.6 filed 21 September 1988.

The oligonucleotide synthesis cycle is performed as follows:

The coupling solution is made up fresh for each step by mixing 6 vol. of 0.5M tetrazole in anhydrous acetonitrile with 5 vol. of a 0.2M solution of the required beta-cyanoethylphosphoramidite. Coupling time is three minutes. Oxidation with a 0.1M solution of  $\text{I}_2$  in  $\text{THF}/\text{pyridine}/\text{H}_2\text{O}$  yields a stable phosphotriester bond. Detritylation of the 5' end with 3% trichloroacetic acid in dichloromethane allows further extension of the oligonucleotide chain. There was no capping step since the excess of phosphoramidites used over reactive sites on the slide was large enough to drive the coupling to completion. After the synthesis is completed, the oligonucleotide is deprotected in 30%  $\text{NH}_3$  for 10 hours at  $55^\circ$ . The chemicals used in the coupling step are moisture-sensitive, and this critical step must be performed under anhydrous conditions in a sealed container, as follows. The shape of the patch

- 16 -

to be synthesised was cut out of a sheet of silicone rubber (76 x 26 x 0.5 mm) which was sandwiched between a microscope slide, derivatised as described above, and a piece of teflon of the same size and thickness. To this was fitted a short piece of plastic tubing that allowed us to inject and withdraw the coupling solution by syringe and to flush the cavity with Argon. The whole assembly was held together by fold-back paper clips. After coupling the set-up was disassembled and the slide put through the subsequent chemical reactions (oxidation with iodine, and detritylation by treatment with TCA) by dipping it into staining jars.

#### EXAMPLE 1.

As a first example we synthesised the sequences oligo-dT<sub>10</sub>-oligo-dT<sub>14</sub> on a slide by gradually decreasing the level of the coupling solution in steps 10 to 14. Thus the 10-mer was synthesised on the upper part of the slide, the 14-mer at the bottom and the 11, 12 and 13-mers were in between. We used 10 pmol oligo-dA<sub>12</sub>, labelled at the 5' end with <sup>32</sup>P by the polynucleotide kinase reaction to a total activity of 1.5 million c.p.m., as a hybridisation probe. Hybridisation was carried out in a perspex (Plexiglas) container made to fit a microscope slide, filled with 1.2 ml of 1M NaCl in TE, 0.1% SDS, for 5 minutes at 20°. After a short rinse in the same solution without oligonucleotide, we were able to detect more than 2000 c.p.s. with a radiation monitor. An autoradiograph showed that all the counts came from the area where the oligonucleotide had been synthesised, i.e. there was no non-specific binding to the glass or to the region that had been derivatised with the linker only. After partial elution in 0.1 M NaCl differential binding to the target is detectable, i.e. less binding to the shorter than the longer oligo-dT. By gradually heating the

- 17 -

slide in the wash solution we determined the  $T_{om}$  (mid-point of transition when 50% eluted) to be  $33^{om}$ . There were no counts detectable after incubation at  $39^o$ .

The hybridisation and melting was repeated eight times with no diminution of the signal. The result is reproducible. We estimate that at least 5% of the input counts were taken up by the slide at each cycle.

#### EXAMPLE 2.

In order to determine whether we would be able to distinguish between matched and mismatched oligonucleotides we synthesised two sequences 3' CCC GCC GCT GGA (cosL) and 3' CCC GCC TCT GGA, which differ by one base at position 7. All bases except the seventh were added in a rectangular patch. At the seventh base, half of the rectangle was exposed in turn to add the two different bases, in two stripes. Hybridisation of cosR probe oligonucleotide (5' GGG CGG CGA CCT) (kinase labelled with  $^{32}P$  to 1.1 million c.p.m., 0.1 M NaCl, TE, 0.1% SDS) was for 5 hours at  $32^o$ . The front of the slide showed 100 c.p.s. after rinsing. Autoradiography showed that annealing occurred only to the part of the slide with the fully complementary oligonucleotide. No signal was detectable on the patch with the mismatched sequence.

#### EXAMPLE 3.

For a further study of the effects of mismatches or shorter sequences on hybridisation behaviour, we constructed two arrays; one (a) of 24 oligonucleotides and the other (b) of 72 oligonucleotides.

These arrays were set out as shown in Table 1(a) and 1(b). The masks used to lay down these arrays were different from those used in previous experiments. Lengths of silicone rubber tubing (1mm c.d.) were glued with silicone rubber cement to the surface of plain microscope slides, in the form of a "U". Clamping

these masks against a derivatised microscope slide produced a cavity into which the coupling solution was introduced through a syringe. In this way only the part of the slide within the cavity came into contact with the phosphoramidite solution. Except in the positions of the mismatched bases, the arrays listed in Table 1 were laid down using a mask which covered most of the width of the slide. Off-setting this mask by 3mm up or down the derivatised slide in subsequent coupling reactions produced the oligonucleotides truncated at the 3' or 5' ends.

For the introduction of mismatches a mask was used which covered half (for array (a)) or one third (for array (b)) of the width of the first mask. The bases at positions six and seven were laid down in two or three longitudinal stripes. This led to the synthesis of oligonucleotides differing by one base on each half (array (a)) or third (array (b)) of the slide. In other positions, the sequences differed from the longest sequence by the absence of bases at the ends.

In array (b), there were two columns of sequences between those shown in Table 1(b), in which the sixth and seventh bases were missing in all positions, because the slide was masked in a stripe by the silicone rubber seal. Thus there were a total of 72 different sequences represented on the slide in 90 different positions.

The 19-mer 5' CTC CTG AGG AGA AGT CTG C was used for hybridisation (2 million cpm, 1.2 ml 0.1M NaCl in TE, 0.1% SDS, 20<sup>o</sup>).

The washing and elution steps were followed by autoradiography. The slide was kept in the washing solution for 5 min at each elution step and then exposed (45 min, intensified). Elution temperatures were 23, 36, 42, 47, 55 and 60<sup>o</sup> C respectively.

As indicated in the table, the oligonucleotides showed different melting behaviour. Short oligonucleotides melted before longer ones, and at 55°C, only the perfectly matched 19-mer was stable, all other oligonucleotides had been eluted. Thus the method can differentiate between a 18-mer and a 19-mer which differ only by the absence of one base at the end. Mismatches at the end of the oligonucleotides and at internal sites can all be melted under conditions where the perfect duplex remains.

Thus we are able to use very stringent hybridisation conditions that eliminate annealing to mismatch sequences or to oligonucleotides differing in length by as little as one base. No other method using hybridisation of oligonucleotides bound to the solid supports is so sensitive to the effects of mismatching.

#### EXAMPLE 4.

To test the application of the invention to diagnosis of inherited diseases, we hybridised the array (a), which carries the oligonucleotide sequences specific for the wild type and the sickle cell mutations of the  $\beta$ -globin gene, with a 110 base pair fragment of DNA amplified from the  $\beta$ -globin gene by means of the polymerase chain reaction (PCR). Total DNA from the blood of a normal individual (1 microgram) was amplified by PCR in the presence of appropriate primer oligonucleotides. The resulting 110 base pair fragment was purified by electrophoresis through an agarose gel. After elution, a small sample (ca. 10 picogram) was labelled by using  $\alpha$ -<sup>32</sup>P-dCTP (50 microCurie) in a second PCR reaction. This PCR contained only the upstream priming oligonucleotide. After 60 cycles of amplification with an extension time of 9 min. the product was removed from precursors by gel filtration. Gel electrophoresis of the radioactive product showed a

- 20 -

major band corresponding in length to the 110 base fragment. One quarter of this product (100,000 c.p.m. in 0.9 M NaCl, TE, 0.1% SDS) was hybridised to the array (a). After 2 hours at 30° ca. 15000 c.p.m. had  
5 been taken up. The melting behaviour of the hybrids was followed as described for the 19-mer in example 3, and it was found that the melting behaviour was similar to that of the oligonucleotide. That is to say, the mismatches considerably reduced the melting temperature  
10 of the hybrids, and conditions were readily found such that the perfectly matched duplex remained whereas the mismatched duplexes had fully melted.

Thus the invention can be used to analyse long fragments of DNA as well oligonucleotides, and this  
15 example shows how it may be used to test nucleic acid sequences for mutations. In particular it shows how it may be applied to the diagnosis of genetic diseases.

#### EXAMPLE 5.

To test an automated system for laying down the  
20 precursors, the cosL oligonucleotide was synthesised with 11 of the 12 bases added in the way described above. For the addition of the seventh base, however, the slide was transferred into an Argon filled chamber containing a pen plotter. The pen of the plotter had  
25 been replaced by a component, fabricated from Nylon, which had the same shape and dimensions as the pen, but which carried a polytetrafluoroethylene (PTFE) tube, through which chemicals could be delivered to the surface of the glass slide which lay on the bed of the  
30 plotter. A microcomputer was used to control the plotter and the syringe pump which delivered the chemicals. The pen, carrying the delivery tube from the syringe, was moved into position above the slide, the pen was lowered and the pump activated to lay down  
35 coupling solution. Filling the pen successively with

- 21 -

6, T and A phosphoramidite solutions an array of twelve spots was laid down in three groups of four, with three different oligonucleotide sequences. After hybridisation to cosR, as described in Example 2, and autoradiography, signal was seen only over the four spots of perfectly matched oligonucleotides, where the dG had been added.

In conclusion, we have demonstrated the following:

1. It is possible to synthesise oligonucleotides in good yield on a flat glass plate.
  2. Multiple sequences can be synthesised on the sample in small spots, at high density, by a simple manual procedure, or automatically using a computer controlled device.
  3. Hybridisation to the oligonucleotides on the plate can be carried out by a very simple procedure. Hybridisation is efficient, and hybrids can be detected by a short autoradiographic exposure.
  4. Hybridisation is specific. There is no detectable signal on areas of the plate where there are no oligonucleotides. We have tested the effects of mismatched bases, and found that a single mismatched base at any position in oligonucleotides ranging in length from 12-mer to 19-mer reduces the stability of the hybrid sufficiently that the signal can be reduced to a very low level, while retaining significant hybridisation to the perfectly matched hybrid.
  5. The oligonucleotides are stably bound to the glass and plates can be used for hybridisation repeatedly.
- The invention thus provides a novel way of analysing nucleotide sequences, which should find a wide range of application. We list a number of potential applications below:

- 22 -

Small arrays of oligonucleotides as fingerprinting  
and mapping tools

Analysis of known mutations including genetic diseases.

5       Example 4 above shows how the invention may be  
used to analyse mutations. There are many  
applications for such a method, including the detection  
of inherited diseases.

Genomic fingerprinting.

10       In the same way as mutations which lead to disease  
can be detected, the method could be used to detect  
point mutations in any stretch of DNA. Sequences are  
now available for a number of regions containing the  
base differences which lead to restriction fragment  
15       length polymorphisms (RFLPs). An array of oligo-  
nucleotides representing such polymorphisms could be  
made from pairs of oligonucleotides representing the  
two allelic restriction sites. Amplification of the  
sequence containing the RFLP, followed by hybridisation  
20       to the plate, would show which alleles were present in  
the test genome. The number of oligonucleotides that  
could be analysed in a single analysis could be quite  
large. Fifty pairs made from selected alleles would  
be enough to give a fingerprint unique to an individual.

Linkage analysis.

25       Applying the method described in the last  
paragraph to a pedigree would pinpoint recombinations.  
Each pair of spots in the array would give the  
information that is seen in the track of the RFLP  
analysis, using gel electrophoresis and blotting, that  
30       is now routinely used for linkage studies. It should  
be possible to analyse many alleles in a single  
analysis, by hybridisation to an array of allelic pairs  
of oligonucleotides, greatly simplifying the methods  
used to find linkage between a DNA polymorphism and  
35       phenotypic marker such as a disease gene.

The examples above could be carried out using the



- 23 -

method we have developed and confirmed by experiments.

Large arrays of oligonucleotides as sequence  
reading tools.

We have shown that oligonucleotides can be  
5 synthesised in small patches in precisely determined  
positions by one of two methods: by delivering the  
precursors through the pen of a pen-plotter, or by  
masking areas with silicone rubber. It is obvious how  
a pen plotter could be adapted to synthesise large  
10 arrays with a different sequence in each position. For  
some applications the array should be a predetermined,  
limited set; for other applications, the array should  
comprise every sequence of a predetermined length. The  
masking method can be used for the latter by laying  
15 down the precursors in a mask which produces  
intersecting lines. There are many ways in which this  
can be done and we give one example for illustration:  
1. The first four bases, A, C, G, T, are laid in four  
broad stripes on a square plate.  
20 2. The second set is laid down in four stripes equal  
in width to the first, and orthogonal to them. The  
array is now composed of all sixteen dinucleotides.  
3. The third and fourth layers are laid down in four  
sets of four stripes one quarter the width of the first  
25 stripes. Each set of four narrow stripes runs within  
one of the broader stripes. The array is now composed  
of all 256 tetranucleotides.  
4. The process is repeated, each time laying down two  
layers with stripes which are one quarter the width of  
30 the previous two layers. Each layer added increases  
the length of the oligonucleotides by one base, and the  
number of different oligonucleotide sequences by a  
factor of four.

The dimensions of such arrays are determined by  
35 the width of the stripes. The narrowest stripe we

have laid is 1mm, but this is clearly not the lowest limit.

There are useful applications for arrays in which part of the sequence is predetermined and part made up of all possible sequences. For example:

#### Characterising mRNA populations.

Most mRNAs in higher eukaryotes have the sequence AAUAAA close to the 3' end. The array used to analyse mRNAs would have this sequence all over the plate. To analyse a mRNA population it would be hybridised to an array composed of all sequences of the type N AATAAAN. For  $m + n = 8$ , which should be enough to give a unique oligonucleotide address to most of the several thousand mRNAs that is estimated to be present in a source such as a mammalian cell, the array would be 256 elements square. The 256 x 256 elements would be laid on the AATAAA using the masking method described above. With stripes of around 1mm, the array would be ca. 256mm square.

This analysis would measure the complexity of the mRNA population and could be used as a basis for comparing populations from different cell types. The advantage of this approach is that the differences in the hybridisation pattern would provide the sequence of oligonucleotides that could be used as probes to isolate all the mRNAs that differed in the populations.

#### Sequence determination.

To extend the idea to determine unknown sequences, using an array composed of all possible oligonucleotides of a chosen length, requires larger arrays than we have synthesised to date. However, it is possible to scale down the size of spot and scale up the numbers to those required by extending the methods we have developed and tested on small arrays. Our experience shows that the method is much simpler in operation than the gel based methods.

TABLE 1

For Examples 3 and 4 array (a) was set out as follows:

|    |    |                           |    |                           |
|----|----|---------------------------|----|---------------------------|
|    | 20 | GAG GAC TCC TCT ACG       | 20 | GAG GAC aCC TCT ACG       |
|    | 36 | GAG GAC TCC TCT GAC G     | 20 | GAC GAC aCC TCT GAC G     |
|    | 36 | GAG GAC TCC TCT AGA CG    | 20 | GAC GAC aCC TCT AGA CG    |
|    | 47 | GAG GAC TCC TCT CAG ACG   | 36 | GAG GAC aCC TCT CAG ACG   |
| 5  | 60 | GAG GAC TCC TCT TCA GAC G | 47 | GAG GAC aCC TCT TCA GAC G |
|    | 56 | .AG GAC TCC TCT TCA GAC G | 42 | .AG GAC aCC TCT TCA GAC G |
|    | 56 | ..G GAC TCC TCT TCA GAC G | 42 | ..G GAC aCC TCT TCA GAC G |
|    | 47 | ... GAC TCC TCT TCA GAC G | 42 | ... GAC aCC TCT TCA GAC G |
|    | 42 | ... .AC TCC TCT TCA GAC G | 36 | ... .AC aCC TCT TCA GAC G |
| 10 | 36 | ... ..C TCC TCT TCA GAC G | 36 | ... ..C aCC TCT TCA GAC G |
|    | 36 | ... ... TCC TCT TCA GAC G | 36 | ... ... aCC TCT TCA GAC G |
|    | 36 | ... ... .CC TCT TCA GAC G | 36 | ... ... .CC TCT TCA GAC G |

For example 3 array (b) was set out as follows:

|    |    |                           |    |                           |    |                           |
|----|----|---------------------------|----|---------------------------|----|---------------------------|
| 15 | 20 | GAG GAt TC                | 20 | GAG GAC TC                | 20 | GAG GAC aC                |
|    | 20 | GAG GAt TCC               | 20 | GAG GAC TCC               | 20 | GAG GAC aCC               |
|    | 20 | GAG GAt TCC T             | 20 | GAG GAC TCC T             | 20 | GAG GAC aCC T             |
|    | 20 | GAG GAt TCC TC            | 20 | GAG GAC TCC TC            | 20 | GAG GAC aCC TC            |
|    | 20 | GAG GAt TCC TCT           | 20 | GAG GAC TCC TCT           | 20 | GAG GAC aCC TCT           |
| 20 | 20 | GAG GAt TCC TCT T         | 20 | GAG GAC TCC TCT T         | 20 | GAG GAC aCC TCT T         |
|    | 20 | GAG GAt TCC TCT TC        | 20 | GAG GAC TCC TCT TC        | 20 | GAG GAC aCC TCT TC        |
|    | 20 | GAG GAt TCC TCT TCA       | 20 | GAG GAC TCC TCT TCA       | 20 | GAG GAC aCC TCT TCA       |
|    | 32 | GAG GAt TCC TCT TCA G     | 42 | GAG GAC TCC TCT TCA G     | 20 | GAG GAC aCC TCT TCA G     |
|    | 32 | GAG GAt TCC TCT TCA GA    | 47 | GAG GAC TCC TCT TCA GA    | 32 | GAG GAC aCC TCT TCA GA    |
| 25 | 42 | GAG GAt TCC TCT TCA GAC   | 52 | GAG GAC TCC TCT TCA GAC   | 42 | GAG GAC aCC TCT TCA GAC   |
|    | 52 | GAG GAt TCC TCT TCA GAC G | 60 | GAG GAC TCC TCT TCA GAC G | 52 | GAG GAC aCC TCT TCA GAC G |
|    | 42 | .AG GAt TCC TCT TCA GAC G | 52 | .AG GAC TCC TCT TCA GAC G | 42 | .AG GAC aCC TCT TCA GAC G |
|    | 42 | ..G GAt TCC TCT TCA GAC G | 52 | ..G GAC TCC TCT TCA GAC G | 42 | ..G GAC aCC TCT TCA GAC G |
|    | 37 | ... GAt TCC TCT TCA GAC G | 47 | ... GAC TCC TCT TCA GAC G | 37 | ... GAC aCC TCT TCA GAC G |
| 30 | 32 | ... .At TCC TCT TCA GAC G | 42 | ... .AC TCC TCT TCA GAC G | 32 | ... .AC aCC TCT TCA GAC G |
|    | 32 | ... ..t TCC TCT TCA GAC G | 42 | ... ..C TCC TCT TCA GAC G | 32 | ... ..C aCC TCT TCA GAC G |
|    | 32 | ... ... TCC TCT TCA GAC G | 32 | ... ... TCC TCT TCA GAC G | 32 | ... ... aCC TCT TCA GAC G |

Between the three columns of array (b) listed above, were two columns, in which bases 6 and 7 from the left were missing in

every line. These sequences all melted at 20 or 32 degrees.

(a,t) mismatch base (.) missing base.

CLAIMS

1. Apparatus for analysing a polynucleotide sequence, comprising a support and attached to a surface thereof an array of the whole or a chosen part of a complete  
5 set of oligonucleotides of chosen lengths, the oligonucleotides being capable of taking part in hybridisation reactions.
2. Apparatus for studying differences between polynucleotide sequences, comprising a support and attached  
10 to a surface thereof an array of the whole or a chosen part of a complete set of oligonucleotides of chosen lengths comprising the polynucleotide sequences, the oligonucleotides being capable of taking part in hybridisation reactions.
- 15 3. Apparatus as claimed in claim 2, wherein the array comprises one or more pairs of oligonucleotides of chosen lengths.
4. Apparatus as claimed in claim 3, wherein the array comprises one or more pairs of oligonucleotides of  
20 chosen lengths representing normal and mutant versions of a point mutation to be studied.
5. Apparatus as claimed in any one of claims 1 to 4, wherein the chosen length is from 8 to 20 nucleotides.
- 25 6. Apparatus as claimed in any one of claims 1 to 5, wherein the surface of the support to which the oligonucleotides are attached is of glass.
7. Apparatus as claimed in any one of claims 1 to 6, wherein each oligonucleotide is bound to the support through a covalent link.
- 30 8. A method of analysing a polynucleotide sequence, by the use of a support to the surface of which is attached an array of the whole or a chosen part of a complete set of oligonucleotides of chosen lengths, which method comprises labelling the polynucleotide  
35 sequence or fragments thereof to form labelled material,

applying the labelled material under hybridisation conditions to the array, and observing the location of the label on the surface associated with particular members of the set of oligonucleotides.

5 9. A method according to claim 8, applied to the study of differences between polynucleotide sequences, wherein the array is of the whole or a chosen part of the complete set of oligonucleotides of chosen lengths comprising the polynucleotide sequences.

10 10. A method as claimed in claim 9, wherein the array comprises one or more pairs of oligonucleotides of chosen lengths.

11. A method as claimed in claim 10, wherein the array comprises one or more pairs of oligonucleotides of  
15 chosen lengths representing normal and mutant versions of a point mutation being studied.

12. A method according to any one of claims 8 to 11, wherein the polynucleotide sequence is randomly degraded to form a mixture of oligomers of a chosen  
20 length, the mixture being thereafter labelled to form the labelled material.

13. A method as claimed in claim 12, wherein the oligomers are labelled with <sup>32</sup>P.

14. A method as claimed in any one of claims 8 to 13,  
25 wherein the chosen length is from 8 to 20 nucleotides.

30

35

# INTERNATIONAL SEARCH REPORT

International Application No **PCT/GB 89/00460**

| <b>I. CLASSIFICATION OF SUBJECT MATTER</b> (If several classification symbols apply, indicate all) <sup>6</sup><br>According to International Patent Classification (IPC) or to both National Classification and IPC<br><b>IPC<sup>4</sup>: C 12 Q 1/68</b>  |  |                                     |   |  |  |  |   |      |          |  |  |          |   |  |
|--|--|-------------------------------------|---|--|--|--|---|------|----------|--|--|----------|---|--|
| <b>II. FIELDS SEARCHED</b><br><div style="text-align: right; font-size: small;">Minimum Documentation Searched <sup>7</sup></div> <table style="width: 100%; border: none;"> <tr> <td style="width: 25%; border: none;">Classification System: --</td> <td style="border: none;">Classification Symbols</td> </tr> <tr> <td style="border: 1px solid black; padding: 5px;"><b>IPC<sup>4</sup></b></td> <td style="border: 1px solid black; padding: 5px;"><b>C 12 Q</b></td> </tr> </table> <div style="text-align: center; font-size: x-small; margin-top: 5px;">Documentation Searched other than Minimum Documentation<br/>to the Extent that such Documents are Included in the Fields Searched <sup>8</sup></div>   |  |                                     | Classification System: --   | Classification Symbols   | <b>IPC<sup>4</sup></b>   | <b>C 12 Q</b>  |   |      |          |  |  |          |   |  |
| Classification System: --  | Classification Symbols   |                                     |   |  |  |  |   |      |          |  |  |          |   |  |
| <b>IPC<sup>4</sup></b>   | <b>C 12 Q</b>  |                                     |   |  |  |  |   |      |          |  |  |          |   |  |
| <b>III. DOCUMENTS CONSIDERED TO BE RELEVANT <sup>9</sup></b> <table border="1" style="width: 100%; border-collapse: collapse; font-size: small;"> <thead> <tr> <th style="width: 10%;">Category <sup>9</sup></th> <th style="width: 60%;">Citation of Document, <sup>11</sup> with Indication, where appropriate, of the relevant passages <sup>12</sup></th> <th style="width: 30%;">Relevant to Claim No. <sup>13</sup></th> </tr> </thead> <tbody> <tr> <td style="text-align: center; vertical-align: top;"><b>X</b></td> <td>GB, A, 2156074 (ORION-YHTYMA OY) 2 October 1985, see the whole document<br/>--</td> <td style="vertical-align: top;">1-14</td> </tr> <tr> <td style="text-align: center; vertical-align: top;"><b>A</b></td> <td>EP, A, 0228075 (MOLECULAR DIAGNOSTIC INC.) 8 July 1987<br/>--</td> <td></td> </tr> <tr> <td style="text-align: center; vertical-align: top;"><b>A</b></td> <td>WO, A, 85/01051 (MOLECULAR BIOSYSTEMS INC.) 14 March 1985<br/>cited in the application<br/>----</td> <td></td> </tr> </tbody> </table>   |  |                                     | Category <sup>9</sup>   | Citation of Document, <sup>11</sup> with Indication, where appropriate, of the relevant passages <sup>12</sup> | Relevant to Claim No. <sup>13</sup>                                | <b>X</b>   | GB, A, 2156074 (ORION-YHTYMA OY) 2 October 1985, see the whole document<br>-- | 1-14 | <b>A</b> | EP, A, 0228075 (MOLECULAR DIAGNOSTIC INC.) 8 July 1987<br>-- |  | <b>A</b> | WO, A, 85/01051 (MOLECULAR BIOSYSTEMS INC.) 14 March 1985<br>cited in the application<br>---- |  |
| Category <sup>9</sup>  | Citation of Document, <sup>11</sup> with Indication, where appropriate, of the relevant passages <sup>12</sup> | Relevant to Claim No. <sup>13</sup> |   |  |  |  |   |      |          |  |  |          |   |  |
| <b>X</b>   | GB, A, 2156074 (ORION-YHTYMA OY) 2 October 1985, see the whole document<br>--                                  | 1-14                                |   |  |  |  |   |      |          |  |  |          |   |  |
| <b>A</b>   | EP, A, 0228075 (MOLECULAR DIAGNOSTIC INC.) 8 July 1987<br>--   |                                     |   |  |  |  |   |      |          |  |  |          |   |  |
| <b>A</b>   | WO, A, 85/01051 (MOLECULAR BIOSYSTEMS INC.) 14 March 1985<br>cited in the application<br>----                  |                                     |   |  |  |  |   |      |          |  |  |          |   |  |
| <div style="display: flex; justify-content: space-between; font-size: x-small;"> <div style="width: 48%;"> <p><sup>10</sup> Special categories of cited documents: <sup>10</sup></p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 48%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"A" document member of the same patent family</p> </div> </div> |  |                                     |   |  |  |  |   |      |          |  |  |          |   |  |
| <b>IV. CERTIFICATION</b> <table style="width: 100%; border: none;"> <tr> <td style="width: 50%; border: none; vertical-align: top;">           Date of the Actual Completion of the International Search<br/> <b>1st August 1989</b> </td> <td style="width: 50%; border: none; vertical-align: top;">           Date of Mailing of this International Search Report<br/> <b>28. 08. 89</b> </td> </tr> <tr> <td style="width: 50%; border: none; vertical-align: top;">           International Searching Authority<br/> <b>EUROPEAN PATENT OFFICE</b> </td> <td style="width: 50%; border: none; vertical-align: top;">           Signature of Authorized Officer<br/> <b>M. VAN MOL</b> </td> </tr> </table>  |  |                                     | Date of the Actual Completion of the International Search<br><b>1st August 1989</b> | Date of Mailing of this International Search Report<br><b>28. 08. 89</b>                                       | International Searching Authority<br><b>EUROPEAN PATENT OFFICE</b> | Signature of Authorized Officer<br><b>M. VAN MOL</b> |   |      |          |  |  |          |   |  |
| Date of the Actual Completion of the International Search<br><b>1st August 1989</b>  | Date of Mailing of this International Search Report<br><b>28. 08. 89</b>                                       |                                     |   |  |  |  |   |      |          |  |  |          |   |  |
| International Searching Authority<br><b>EUROPEAN PATENT OFFICE</b>   | Signature of Authorized Officer<br><b>M. VAN MOL</b>   |                                     |   |  |  |  |   |      |          |  |  |          |   |  |

**ANNEX TO THE INTERNATIONAL SEARCH REPORT  
ON INTERNATIONAL PATENT APPLICATION NO.**

GB 8900460

SA 28452

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on 21/08/89. The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

| Patent document<br>cited in search report | Publication<br>date | Patent family<br>member(s) | Publication<br>date |
|---|---------------------|----------------------------|---------------------|
| GB-A- 2156074                             | 02-10-85            | AU-B- 577568               | 29-09-88            |
|   |                     | AU-A- 3842285              | 22-08-85            |
|   |                     | BE-A- 901671               | 29-05-85            |
|   |                     | CA-A- 1248895              | 17-01-89            |
|   |                     | DE-A- 3505287              | 05-09-85            |
|   |                     | FR-A- 2559783              | 23-08-85            |
|   |                     | JP-A- 60188100             | 25-09-85            |
|   |                     | LU-A- 85768                | 24-07-85            |
|   |                     | NL-A- 8500424              | 16-09-85            |
|   |                     | SE-A- 8500733              | 18-08-85            |
|   |                     | US-A- 4731325              | 15-03-88            |
| EP-A- 0228075                             | 08-07-87            | JP-A- 62228300             | 07-10-87            |
| WO-A- 8501051                             | 14-03-85            | AU-B- 575586               | 04-08-88            |
|   |                     | AU-A- 3319684              | 29-03-85            |
|   |                     | EP-A- 0155950              | 02-10-85            |
|   |                     | JP-T- 60502155             | 12-12-85            |